

Superconductivity in the Model with non Cooper Pairs

I.M.Yurin, V.B.Kalinin

*Institute of Physical Chemistry, Leninskiy prosp. 31, GSP-1, Moscow, 119991, Russia**

On a basis of earlier substantiated expression for effective potential of electron-electron attraction in metals the assumption of an opportunity of formation of classically bound pairs is put forward. It was shown that in distinction from the Cooper pair, the energy of the electron pair in this approach is negative in the centrum of mass of the pair. This fact changes the pattern of the superconductivity phenomenon in the proposed model. First, the gap in the one-particle spectrum appears due to the effect of a "mean field" on the energy of electron state from the side of occupied states, the nearest neighbors over the momenta grid. Second, the condensate is formed by electron states with energies below that of the gap edge. No Bose condensation does occur in the system proceeding from the proposed model. In quasi-classical approximation Londons' hypothesis is substantiated.

I. INTRODUCTION

At present, it is the superconductivity theory put forward by Bardeen, Cooper and Schrieffer in 1957 that is considered generally accepted ¹. In the framework of this theory, one succeeded in clarifying a great many of experiments devoted to investigating so-called conventional superconductors, i.e., those with rather low transition temperatures. At the time of discovering high-temperature superconductivity (HTSC), the most optimistic assessments of BCS theory yielded the value of about 40 K as the possible maximal transition temperature T_S .

The discovery of HTSC divided research workers into two big groups. One part of research workers is continuing to consider the BCS theory to be suitable for describing HTSC: for this purpose, it would be sufficient to substantiate the introduction of large values of constants of electron-phonon interaction. Now, other research workers consider the BCS theory to be applicable to conventional superconductivity. However, for example, in cuprates one has to search for novel mechanisms of the pairing of electrons. As concerns the authors of the present paper, they are in obvious minority, forming a group considering it to be necessary to radically revise the existing superconductivity theory. In the framework of the novel theory that is being developed in our works the HTSC has been naturally substantiated, whereas the descriptions of superconductivity and HTSC are lacking essential differences.

In order to properly understand the present-day state of the superconductivity theory, it is needed to thoroughly consider the generally accepted BCS theory. The BCS Hamiltonian binds the pairs with zero momenta only, and has the following form:

$$H_{BCS} = \sum_{\sigma, \mathbf{k}} T_{\mathbf{k}} c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}} - \frac{g}{\Omega} \sum_{\substack{E_F - \omega_D < T_{\mathbf{k}} < E_F + \omega_D \\ E_F - \omega_D < T_{\mathbf{p}} < E_F + \omega_D}} c_{\uparrow, \mathbf{p}}^+ c_{\downarrow, -\mathbf{p}}^+ c_{\downarrow, -\mathbf{k}} c_{\uparrow, \mathbf{k}}, \quad (1)$$

where $T_{\mathbf{k}}$ is the kinetic energy, $c_{\sigma, \mathbf{k}}^+$ and $c_{\sigma, \mathbf{k}}$ are the operators of the creation and annihilation of the electron with momentum $\mathbf{k}$ and spin index $\sigma = \pm \frac{1}{2}$ (or $\sigma = \uparrow, \downarrow$ respectively), E_F is the Fermi energy, Ω is the volume of the system, and ω_D is the Debye frequency.

A rigorous analysis shows that Hamiltonian (1) describes a dielectric rather than a superconductor. It is possible to become convinced in that already by scrutinizing so-called "elementary Cooper superconductor", which is essentially a single pair of electrons bound together by the interaction term of the Hamiltonian (1) (it is curious that in distinction from the classically bound pairs, the energy of Cooper pair in the system of the centrum of masses with realistically chosen parameters of the model is positive).

Let us assume for the sake of simplicity that $\omega_D = E_F$. If we apply to the Cooper superconductor a voltage V of such a value that the condition $2eV < E_b$ is fulfilled, where E_b is the energy of a pair to be counted from the bottom of the conductivity zone and e is the electron charge, then the response of such a system to the electrical field will be calculated in the framework of the perturbation theory, for the energy of the Cooper pair is separated from all other states by an energy gap E_b . Thus, the Cooper "superconductor" is actually incapable of screening the electrical field, that is, in the BCS system even the properties of normal metal have been lost.

It may seem that Bose condensation for many pairs can substantially change the situation. This supposition refutes the strict solution of a many electron problem for the reduced BCS Hamiltonian, as given in study ². It turns out that the ground state of such a system is also separated from other states by an energy gap. Therefore, in order to induce current, it is necessary to consume a finite quantity of energy, which is incompatible with the claimed superconductivity properties. In other words, the BCS model actually describes the Mott hypothetical transition, when the ground state of the system proves to be dielectric, and has nothing in common with the superconductivity problem.

It may seem that we have to deal with a shortcoming of the original formulation of BCS theory. In the further course, when we pass over to considering the bound pairs with the momenta that are different from zero, this shortcoming disappears. Let us show this does not correspond to reality.

So, in the framework of Gor'kov transformation ³, some of the interaction terms of the Hamiltonian

$$H = \sum_{\sigma, \mathbf{k}} T_{\mathbf{k}} c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}} + \sum_{\sigma, \nu} \sum_{\mathbf{p}, \mathbf{k}, \mathbf{q}} V_{\mathbf{q}}^{real} c_{\sigma, \mathbf{p}+\mathbf{q}}^+ c_{\nu, \mathbf{k}-\mathbf{q}}^+ c_{\nu, \mathbf{k}} c_{\sigma, \mathbf{p}} \quad (2)$$

are diagonalized by a transition to the operators of the creation and annihilation of bogolons:

$$\begin{aligned} \gamma_{\sigma, \mathbf{p}}^+ &= u_p c_{\sigma, \mathbf{p}}^+ - 2\sigma v_p c_{-\sigma, -\mathbf{p}}, \\ \gamma_{\sigma, \mathbf{p}} &= u_p c_{\sigma, \mathbf{p}} - 2\sigma v_p c_{-\sigma, -\mathbf{p}}^+. \end{aligned} \quad (3)$$

The parameters u_p and v_p are determined in the framework of the self-consistent Hartree–Fock procedure based on equation for the bogolons' spectrum $[H, \gamma_{\sigma, \mathbf{p}}^+] = E_p \gamma_{\sigma, \mathbf{p}}^+$. The appearance of the gap in the spectrum is treated as a transition into the superconductivity state.

Now we are going to note that equation $[H, \gamma_{\sigma, \mathbf{p}}^+] = E_p \gamma_{\sigma, \mathbf{p}}^+$ has a physical sense only in the case where the vacuum state $|BCS\rangle$, satisfying the condition $\gamma_{\sigma, \mathbf{p}} |BCS\rangle = 0$, is stationary (in the BCS theory this vacuum state is considered the ground state of the system). For further purpose, it is sufficient to write the Hamiltonian (2) in the operators $\gamma_{\sigma, \mathbf{p}}^+$ and $\gamma_{\sigma, \mathbf{p}}$.

In the case of the BCS Hamiltonian accounting for the terms with the structure $\gamma^+ \gamma^+ \gamma^+ \gamma^+$ we derive the following expression $|\delta\psi| \sim N/\Omega$, where $\delta\psi$ is a correction of the first order of the perturbation theory to the wave function of the state $|BCS\rangle$, and N is the mean number of electrons in the system. But, in a general case for Hamiltonian (2), the analogous terms give $|\delta\psi| \sim N^{3/2}/\Omega$, or $|\delta\psi| \sim n\sqrt{N}$, where n is concentration of electrons in the system. Therefore, the Gor'kov procedure is definitely incorrect when describing real systems with large dimensions.

Now, let us consider Eliashberg's equations ⁴, which are frequently referred to as the modern formulation of BCS theory ⁵. These equations were derived from the Fröhlich model in 1960. At the same time, it is long known ⁶ that in the Fröhlich model it is impracticable to effect the consistent taking into account of the Coulomb electron-electron interaction in the procedure of screening electron-phonon potential. This circumstance reveals the irrelevance of the Eliashberg's equations for Green's functions of electrons and phonons with respect to the essential point of the renormalization problem. Moreover, the use of anomalous expectation values makes the determination of the ground state of the system as unsubstantiated as in the case of Gor'kov transformation.

The consistent renormalization in the system of electrons and phonons was for the first time carried into effect in study ⁷. It has been shown that introduction of an additional kinetic terms $\sim q^{-1} (2b_{\mathbf{q}}^+ b_{\mathbf{q}} - b_{\mathbf{q}}^+ b_{-\mathbf{q}}^+ - b_{\mathbf{q}} b_{-\mathbf{q}})$, where $b_{\mathbf{q}}^+$ and $b_{\mathbf{q}}$ are the operators of the creation and annihilation of the phonons of longitudinal polarization with momentum $\mathbf{q}$, enables one to balance a series of singularities in the long-wave limit, that arise in the process of renormalization.

The consideration of the long-wave limit of the electron-phonon and, accordingly, electron-electron interaction is especially relevant in studying the electron pairs that are bound in the classical sense. In fact, it is easy to take into account the short-wave interaction in the framework of the perturbation theory because of high values of the arising energy denominators. Therefore, in distinction from the BCS theory, it is just the long-wave interaction that should play the crucial role in the substantiation of physically relevant superconductivity theories.

In the framework of the same works ⁷, when taking into consideration electronic correlations, we have derived an expression for the effective Hamiltonian of degenerated Fermi gas in the model of monatomic metal:

$$H_{eff} = \sum_{\sigma, \mathbf{k}} T_{\mathbf{k}} c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}} + \sum_{\sigma, \nu} \sum_{\mathbf{p}, \mathbf{k}, \mathbf{q}} \tilde{U}_{\mathbf{q}}^{\mathbf{p}, \mathbf{k}} c_{\sigma, \mathbf{p}+\mathbf{q}}^+ c_{\nu, \mathbf{k}-\mathbf{q}}^+ c_{\nu, \mathbf{k}} c_{\sigma, \mathbf{p}}. \quad (4)$$

In that, with $p, k \gg mS$, we have

$$\tilde{U}_{\mathbf{q}}^{\mathbf{p}, \mathbf{k}} \sim - \left(\frac{zm}{3M} \right)^2 \frac{e^2}{\varepsilon q^2} \frac{K_F^2}{q^2 - 4m^2 s^2} \frac{K_F^2}{q^2 - 4m^2 s_1^2}, \quad (5)$$

where z is the number of conductivity electrons per cell, m is the zone mass of electron, M is the mass of ion, ε is the dielectric permittivity of the electrons of valence zone, K_F is the Fermi wave vector, s is the initial velocity of phonons, which is linked with the observed sound velocity S in metal by a relationship $S = \sqrt{s^2 + \frac{zm}{3M} V_F^2}$, V_F is the Fermi velocity, $s_1^2 = \left(1 + \frac{zm}{3M} \frac{\varepsilon K_F^2}{\lambda^2} \right) s^2$, and λ^{-1} is the Thomas-Fermi screening length. In study ⁷, we actually considered a doped semiconductor. In such a case, the electron states may be divided into the states of valence and conductivity zones. In this, the values of dielectric permittivity ε and the initial velocity of phonons s are of explicit

parent dielectric origin. Future investigations will certainly show how correct is the approximation suggested above for describing the metals of periodic system.

Now, we are going to make several remarks pertaining to the subject-matter of this paper. Firstly, the presence of singularities in Eq. (5) is not at all unavoidable in the framework of the approach being developed. As a matter of fact, taking into account the phonon and electron scattering will be conducive to the disappearance of singularities in Eq. (5): this may be achieved by introducing supplementary terms to the initial Hamiltonian introduced in ⁷.

Secondly, one has to perceive distinctly the peculiarities of the construction of the ground state in the framework of the proposed approach. From the very beginning, we believe the ground state of the system may be represented in the following form:

$$|\Phi_0\rangle = \prod_{\sigma, k < K_F} C_{\sigma, \mathbf{k}}^+ |0\rangle, \quad (6)$$

where $C_{\sigma, \mathbf{k}}^+$ are the operators of the creation of quasi-particles originating from electrons.

During the renormalization procedure, taking into account electron correlations, and solving other problems, these operators are refined; however, even the appearance of the bound states of Hamiltonian H_{eff} does not change equation (6). This circumstance qualitatively distinguishes our approach from the most studies, in which the Hamiltonians presupposing the presence of bound states are considered to be initial ones, and, accordingly, the determining of the ground state of the system represents a separate problem.

II. CONSTRUCTION OF THE MODEL HAMILTONIAN

The effective electron-electron interaction (4) was obtained in the first order of the perturbation theory with respect to the small parameter $\tilde{U}/T$. Let us try to generalize the obtained results for the case where the perturbation theory does not work. So, after the completion of the renormalization procedure, let us consider the following Hamiltonian:

$$H = \sum_{\sigma, \mathbf{k}} T_{\mathbf{k}} c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}} + \sum_{\sigma, \nu} \sum_{\mathbf{p}, \mathbf{k}, \mathbf{q}} U_{\mathbf{q}}^{\mathbf{p}, \mathbf{k}} c_{\sigma, \mathbf{p}+\mathbf{q}}^+ c_{\nu, \mathbf{k}-\mathbf{q}}^+ c_{\nu, \mathbf{k}} c_{\sigma, \mathbf{p}}. \quad (7)$$

The scheme of introducing the concept of effective Hamiltonian is similar to that set forth in papers ⁷. At first, we consider the Hamiltonian H_{eff} that, as one would think, is even formally not linked with Eq. (7):

$$\begin{aligned} H_{eff} = & \sum_{\sigma, \mathbf{k}} T_{\mathbf{k}} c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}} \\ & + \frac{1}{2} \sum_{\mathbf{p}, \mathbf{k}, \mathbf{q}} \tilde{U}_{0, \mathbf{q}}^{\mathbf{p}, \mathbf{k}} \left(c_{\uparrow, \mathbf{p}+\mathbf{q}}^+ c_{\downarrow, \mathbf{k}-\mathbf{q}}^+ - c_{\downarrow, \mathbf{p}+\mathbf{q}}^+ c_{\uparrow, \mathbf{k}-\mathbf{q}}^+ \right) (c_{\downarrow, \mathbf{k}} c_{\uparrow, \mathbf{p}} - c_{\uparrow, \mathbf{k}} c_{\downarrow, \mathbf{p}}) \\ & + \sum_{\sigma} \sum_{\mathbf{p}, \mathbf{k}, \mathbf{q}} \tilde{U}_{1, \mathbf{q}}^{\mathbf{p}, \mathbf{k}} c_{\sigma, \mathbf{p}+\mathbf{q}}^+ c_{\sigma, \mathbf{k}-\mathbf{q}}^+ c_{\sigma, \mathbf{k}} c_{\sigma, \mathbf{p}} \\ & + \frac{1}{2} \sum_{\mathbf{p}, \mathbf{k}, \mathbf{q}} \tilde{U}_{1, \mathbf{q}}^{\mathbf{p}, \mathbf{k}} \left(c_{\uparrow, \mathbf{p}+\mathbf{q}}^+ c_{\downarrow, \mathbf{k}-\mathbf{q}}^+ + c_{\downarrow, \mathbf{p}+\mathbf{q}}^+ c_{\uparrow, \mathbf{k}-\mathbf{q}}^+ \right) (c_{\downarrow, \mathbf{k}} c_{\uparrow, \mathbf{p}} + c_{\uparrow, \mathbf{k}} c_{\downarrow, \mathbf{p}}). \end{aligned} \quad (8)$$

Using transformation

$$\begin{aligned} C_{\sigma, \mathbf{p}}^+ &= c_{\sigma, \mathbf{p}}^+ + \sum_{\nu} \sum_{\mathbf{k}, \mathbf{q}} \theta_{\sigma, \nu, \mathbf{q}}^{\mathbf{p}, \mathbf{k}} c_{\sigma, \mathbf{p}+\mathbf{q}}^+ c_{\nu, \mathbf{k}-\mathbf{q}}^+ c_{\nu, \mathbf{k}}, \\ C_{\sigma, \mathbf{p}} &= c_{\sigma, \mathbf{p}} + \sum_{\nu} \sum_{\mathbf{k}, \mathbf{q}} \theta_{\sigma, \nu, \mathbf{q}}^{* \mathbf{p}, \mathbf{k}} c_{\nu, \mathbf{k}} c_{\nu, \mathbf{k}-\mathbf{q}} c_{\sigma, \mathbf{p}+\mathbf{q}}, \end{aligned} \quad (9)$$

the Hamiltonian H_{eff} is transformed to the following form

$$\begin{aligned} H_{eff} = & \sum_{\sigma, \mathbf{k}} T_{\mathbf{k}} C_{\sigma, \mathbf{k}}^+ C_{\sigma, \mathbf{k}} \\ & + \frac{1}{2} \sum_{\sigma, \mathbf{p} \neq \mathbf{k}} E_1^{\mathbf{p}, \mathbf{k}} C_{\sigma, \mathbf{p}}^+ C_{\sigma, \mathbf{k}}^+ C_{\sigma, \mathbf{k}} C_{\sigma, \mathbf{p}} \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{4} \sum_{\mathbf{p} \neq \mathbf{k}} E_1^{\mathbf{p},\mathbf{k}} \left(C_{\uparrow,\mathbf{p}}^+ C_{\downarrow,\mathbf{k}}^+ + C_{\downarrow,\mathbf{p}}^+ C_{\uparrow,\mathbf{k}}^+ \right) (C_{\downarrow,\mathbf{k}} C_{\uparrow,\mathbf{p}} + C_{\uparrow,\mathbf{k}} C_{\downarrow,\mathbf{p}}) \\
& + \sum_{\mathbf{p}} E_0^{\mathbf{p},\mathbf{p}} C_{\uparrow,\mathbf{p}}^+ C_{\downarrow,\mathbf{p}}^+ C_{\downarrow,\mathbf{p}} C_{\uparrow,\mathbf{p}} \\
& + \frac{1}{4} \sum_{\mathbf{p} \neq \mathbf{k}} E_0^{\mathbf{p},\mathbf{k}} \left(C_{\uparrow,\mathbf{p}}^+ C_{\downarrow,\mathbf{k}}^+ - C_{\downarrow,\mathbf{p}}^+ C_{\uparrow,\mathbf{k}}^+ \right) (C_{\downarrow,\mathbf{k}} C_{\uparrow,\mathbf{p}} - C_{\uparrow,\mathbf{k}} C_{\downarrow,\mathbf{p}}).
\end{aligned} \tag{10}$$

In distinction from study ⁷, in Eq. (8) it has been taken into account that in higher than the first orders of the perturbation theory with respect to parameter the effective potentials for singlet and triplet pairs may differ from each other. Moreover, the parameters of transformation (9) will be determined, not from the perturbation theory, but will be reassigned by the expressions of a more general form:

$$\begin{aligned}
\theta_{\sigma,\sigma,\mathbf{q}}^{\mathbf{p},\mathbf{k}} &= \frac{1}{2} \left(\delta_{\mathbf{k}-\mathbf{p}}^{\mathbf{q}} - \delta_{\mathbf{q}}^0 \right) + \chi_{1,\mathbf{q}}^{\mathbf{p},\mathbf{k}}, \\
\theta_{\sigma,-\sigma,\mathbf{q}}^{\mathbf{p},\mathbf{k}} &= -\delta_{\mathbf{q}}^0 + \chi_{0,\mathbf{q}}^{\mathbf{p},\mathbf{k}} + \chi_{1,\mathbf{q}}^{\mathbf{p},\mathbf{k}},
\end{aligned} \tag{11}$$

where $\delta_{\dots}$ is a Kronecker's 3D-symbol, $\chi_{0,\mathbf{q}}^{\mathbf{p},\mathbf{k}}$ and $\chi_{1,\mathbf{q}}^{\mathbf{p},\mathbf{k}}$ are defined by equations on the two-particle eigen states of the Hamiltonian H_{eff} (see Eq. (8) and Appendix):

$$\begin{aligned}
H_{eff} \left| S_1^{\mathbf{p},\mathbf{k}} \right\rangle &= \left(T_{\mathbf{p}} + T_{\mathbf{k}} + E_1^{\mathbf{p},\mathbf{k}} \right) \left| S_1^{\mathbf{p},\mathbf{k}} \right\rangle, \\
H_{eff} \left| S_0^{\mathbf{p},\mathbf{k}} \right\rangle &= \left(T_{\mathbf{p}} + T_{\mathbf{k}} + E_0^{\mathbf{p},\mathbf{k}} \right) \left| S_0^{\mathbf{p},\mathbf{k}} \right\rangle,
\end{aligned} \tag{12}$$

where $\left| S_1^{\mathbf{p},\mathbf{k}} \right\rangle = \sum_{\mathbf{q}} \chi_{1,\mathbf{q}}^{\mathbf{p},\mathbf{k}} c_{\uparrow,\mathbf{p}+\mathbf{q}}^+ c_{\uparrow,\mathbf{k}-\mathbf{q}}^+ |0\rangle$ and $\left| S_0^{\mathbf{p},\mathbf{k}} \right\rangle = \sum_{\mathbf{q}} \chi_{0,\mathbf{q}}^{\mathbf{p},\mathbf{k}} \left(c_{\uparrow,\mathbf{p}+\mathbf{q}}^+ c_{\downarrow,\mathbf{k}-\mathbf{q}}^+ - c_{\downarrow,\mathbf{p}+\mathbf{q}}^+ c_{\uparrow,\mathbf{k}-\mathbf{q}}^+ \right) |0\rangle$.

The equations on the effective potentials $\tilde{U}_{i,\mathbf{q}}^{\mathbf{p},\mathbf{k}}$ are derived in the following manner. Formally, the transformation is introduced, which is reciprocal to transformation (9):

$$\begin{aligned}
c_{\sigma,\mathbf{p}}^+ &= C_{\sigma,\mathbf{p}}^+ + \sum_{\nu} \sum_{\mathbf{k},\mathbf{q}} \theta_{\sigma,\nu,\mathbf{q}}^{*\mathbf{k}-\mathbf{q},\mathbf{p}+\mathbf{q}} C_{\sigma,\mathbf{p}+\mathbf{q}}^+ C_{\nu,\mathbf{k}-\mathbf{q}}^+ C_{\nu,\mathbf{k}}, \\
c_{\sigma,\mathbf{p}} &= C_{\sigma,\mathbf{p}} + \sum_{\nu} \sum_{\mathbf{k},\mathbf{q}} \theta_{\sigma,\nu,\mathbf{q}}^{\mathbf{k}-\mathbf{q},\mathbf{p}+\mathbf{q}} C_{\nu,\mathbf{k}}^+ C_{\nu,\mathbf{k}-\mathbf{q}} C_{\sigma,\mathbf{p}+\mathbf{q}}.
\end{aligned} \tag{13}$$

Using Eq. (13), the Hamiltonian (7) is written in operators C^+ and C . In such a written form, the terms appear in the Hamiltonian having the structure $C^+ C^+ C^+ C C C$, which, in the random phase approximation can be reduced to the form $\langle C^+ C \rangle C^+ C^+ C C$ (here and further the broken brackets $\langle \dots \rangle$ designate the averaging with respect to the ground state). These terms and some other arising terms with the structure $C^+ C^+ C C$ define supplementary potentials $\tilde{D}_{0,\mathbf{q}}^{\mathbf{p},\mathbf{k}}$ and $\tilde{D}_{1,\mathbf{q}}^{\mathbf{p},\mathbf{k}}$. Then the self-consistency equations on the effective potentials $\tilde{U}_{i,\mathbf{q}}^{\mathbf{p},\mathbf{k}}$ become

$$\tilde{U}_{i,\mathbf{q}}^{\mathbf{p},\mathbf{k}} = U_{\mathbf{q}}^{\mathbf{p},\mathbf{k}} + \tilde{D}_{i,\mathbf{q}}^{\mathbf{p},\mathbf{k}} \tag{14}$$

In the case where the perturbation theory is applicable, i.e.,

$$\begin{aligned}
\chi_{0,\mathbf{q}}^{\mathbf{p},\mathbf{k}} &\approx \frac{1}{2} \left(\delta_{\mathbf{q}}^0 + \delta_{\mathbf{k}-\mathbf{p}}^{\mathbf{q}} \right), \\
\chi_{1,\mathbf{q}}^{\mathbf{p},\mathbf{k}} &\approx \frac{1}{2} \left(\delta_{\mathbf{q}}^0 - \delta_{\mathbf{k}-\mathbf{p}}^{\mathbf{q}} \right),
\end{aligned} \tag{15}$$

Eqs. (14) little differ from those derived in study⁷. We are now interested in the case, where some solutions form the states of discrete spectrum, and relationships (15) corresponding to these states are not fulfilled.

Let us restrict ourselves to the consideration of the case where the Hamiltonian has one bound-state solution with a zero spin for pairs with momentum $\mathbf{P}$. For all the remaining unbound pairs with $\mathbf{p} + \mathbf{k} = \mathbf{P}$, the following equation is fulfilled: $\chi_{0,\mathbf{q}}^{\mathbf{p},\mathbf{k}} \approx \frac{1}{2} \left(\delta_{\mathbf{q}}^0 + \delta_{\mathbf{k}-\mathbf{p}}^{\mathbf{q}} \right)$. Determine for which pair this relationship has not been fulfilled (it is the pair forming the discrete spectrum).

For this purpose, note once more that it is the long-wave part of the electron-electron interaction, which plays the main role in the formation of bound states. When considering the electron-electron interaction of two-particle states

in the framework of the perturbation theory, the energy denominator $\sim (\mathbf{p} - \mathbf{k})\mathbf{q}$. Hence, it attains the minimal value when $\mathbf{p} \approx \mathbf{k}$. That is why the bound pair originates from the state of two free electrons that are located on one site of the momentum grid or on the neighboring ones.

We will reckon that the pair binding energy E_b considerably exceeds corrections $E_i^{\mathbf{p},\mathbf{k}}$ for the pairs whose spectrum is well described by the perturbation theory. Then, for a crystal with dimensions $L \times L \times L$ and periodic boundary conditions, we arrive to the following model Hamiltonian:

$$\begin{aligned} H_{mod} = & \sum_{\sigma,\mathbf{k}} T_{\mathbf{k}} C_{\sigma,\mathbf{k}}^+ C_{\sigma,\mathbf{k}} \\ & - \sum_{\mathbf{k}} E_b C_{\uparrow,\mathbf{k}}^+ C_{\downarrow,\mathbf{k}}^+ C_{\downarrow,\mathbf{k}} C_{\uparrow,\mathbf{k}} \\ & - \frac{1}{4} \sum E_b \left(C_{\uparrow,\mathbf{p}}^+ C_{\downarrow,\mathbf{k}}^+ - C_{\downarrow,\mathbf{p}}^+ C_{\uparrow,\mathbf{k}}^+ \right) (C_{\downarrow,\mathbf{k}} C_{\uparrow,\mathbf{p}} - C_{\uparrow,\mathbf{k}} C_{\downarrow,\mathbf{p}}) \\ & \{ \mathbf{p} \neq \mathbf{k}; |\mathbf{p}_\alpha - \mathbf{k}_\alpha| \leq 2\pi/L, \alpha = x, y, z \}. \end{aligned} \quad (16)$$

The energy of a quasi-particle $\tilde{E}_{\mathbf{p}}$ in the mean field approximation accounting for the nearest neighbors over the momenta grid is determined by the following equation:

$$\tilde{E}_{\mathbf{p}} C_{\sigma,\mathbf{k}}^+ = \left[H_{mod}, C_{\sigma,\mathbf{k}}^+ \right] = (T_{\mathbf{k}} - 14E_b \rho_{\mathbf{k}}) C_{\sigma,\mathbf{k}}^+, \quad (17)$$

where $\rho_{\mathbf{k}}$ is the mean value of the occupation number in the vicinity of the site $\mathbf{k}$.

Eq. (17) enables one to obtain the qualitative picture of changes in the electron spectrum during variation in temperature⁸. At the lowest temperatures, the spectrum has a gap of the magnitude $2\Delta_0 = 14E_b$. As temperature rises, the gap decreases. At the transition point, the spectrum reveals a singularity, whereas at a temperature which is higher than the transition point, the gap completely disappears.

The transition temperature can be estimated from an equation $\partial\rho/\partial E|_{\tilde{E}=\mu} = -\infty$, where μ is the chemical potential of electrons. Assuming for ρ the Fermi–Dirac distribution function, we obtain for the transition temperature T_s :

$$T_s = \frac{\Delta_0}{2}. \quad (18)$$

In distinction from the BCS model, the gap forming in the system is evidently superconducting. It follows from the following speculations: firstly, it is possible to obtain the current state of the system by shifting the wave function of the ground state in the momenta space, without consuming the finite energy; secondly, the collision integral in such a system is frozen out according to the exponential law.

The superconducting condensate forms the ground state of the system. The thermally excited electrons outside the momenta space, which is occupied by condensate, and the holes inside the condensate form normal gas that, contrary to the BCS theory, is electrically neutral. All the processes of scattering in the system occur only for the normal gas. Therefore, it proves to be extremely difficult to estimate the condensate relaxation time with non-zero current. However, it is most likely that here we have to do with astronomic values.

Winding up the description of the procedure of introducing the model Hamiltonian, we note once more that in the case where two-particle solutions of discrete spectrum appear in the system, the transformations (9) and (13) are of formal, auxiliary character for the determination of the model Hamiltonian in the space of wave functions of eigen states of the Hamiltonian H_{eff} . These transformations allow one to derive also expressions for the operators of the values observed, corresponding to the Hamiltonian H_{mod} . Thus, for the operator $c_{\sigma,\mathbf{p}}^+ c_{\sigma,\mathbf{p}+1}$, to be applied to determining the current density, we have the following expression:

$$\begin{aligned} c_{\sigma,\mathbf{p}}^+ c_{\sigma,\mathbf{p}+1} = & C_{\sigma,\mathbf{p}}^+ C_{\sigma,\mathbf{p}+1} \\ & - \sum_{\nu,\mathbf{k}} C_{\sigma,\mathbf{p}}^+ C_{\nu,\mathbf{k}}^+ C_{\nu,\mathbf{k}} C_{\sigma,\mathbf{p}+1} \\ & + \sum_{\nu,\mathbf{k}} \sum_{\mathbf{z},\mathbf{y}} \chi_{1,\mathbf{z}+\mathbf{y}}^{*\mathbf{k},\mathbf{p}} \chi_{1,\mathbf{y}}^{\mathbf{k}+\mathbf{z},\mathbf{p}-\mathbf{z}+1} C_{\sigma,\mathbf{p}}^+ C_{\nu,\mathbf{k}}^+ C_{\nu,\mathbf{k}+\mathbf{z}} C_{\sigma,\mathbf{p}-\mathbf{z}+1} \\ & + \sum_{\nu,\mathbf{k}} \sum_{\mathbf{z},\mathbf{y}} \chi_{0,\mathbf{z}+\mathbf{y}}^{*\mathbf{k},\mathbf{p}} \chi_{0,\mathbf{y}}^{\mathbf{k}+\mathbf{z},\mathbf{p}-\mathbf{z}+1} C_{\sigma,\mathbf{p}}^+ C_{-\sigma,\mathbf{k}}^+ C_{-\sigma,\mathbf{k}+\mathbf{z}} C_{\sigma,\mathbf{p}-\mathbf{z}+1}. \end{aligned} \quad (19)$$

We shall employ this equation to calculate the response of the condensate to electromagnetic field.

III. THE BEHAVIOR OF CONDENSATE IN ELECTROMAGNETIC FIELD

The appearance of electromagnetic field in the system under consideration is described by introducing a supplementary term H_{em} . In this, the total Hamiltonian of the system has the following form:

$$H_{tot} = H_{mod} + H_{em}, \quad (20)$$

where

$$\begin{aligned} H_{em} &= -\frac{1}{c} \sum_{\mathbf{q}} \mathbf{A}_{-\mathbf{q},t} \mathbf{j}_{\mathbf{q}}, \\ \mathbf{A}_{\mathbf{q},t} &= L^{-3/2} \int \mathbf{A}(\mathbf{r}, t) e^{-i\mathbf{q}\mathbf{r}} d\mathbf{r}, \\ \mathbf{j}_{\mathbf{q}} &= -\frac{eL^{-3/2}}{2m} \sum_{\sigma, \mathbf{k}} (2\mathbf{k} + \mathbf{q}) c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}+\mathbf{q}} - \frac{e^2 L^{-3}}{mc} \sum_{\sigma, \mathbf{k}, \mathbf{l}} \mathbf{A}_{\mathbf{q}-\mathbf{l},t} c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}+\mathbf{l}}, \end{aligned} \quad (21)$$

while the expressions for $c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}+\mathbf{q}}$ and $c_{\sigma, \mathbf{k}}^+ c_{\sigma, \mathbf{k}+\mathbf{l}}$ may be substituted from Eq. (19).

We will search for the wave function of the condensate coinciding the wave function of the ground state of the system in the following form:

$$|\Phi_t\rangle = \prod_{\sigma, \mathbf{k} < K_F} \tilde{C}_{\sigma, \mathbf{k}}^+(t) |0\rangle, \quad (22)$$

whereas for $\tilde{C}_{\sigma, \mathbf{k}}^+(t)$ we assume the fulfillment of the following relationship:

$$i \frac{\partial \tilde{C}_{\sigma, \mathbf{k}}^+(t)}{\partial t} = [H_{tot}, \tilde{C}_{\sigma, \mathbf{k}}^+(t)]. \quad (23)$$

It is presumably obvious that Schrödinger's equation $i \frac{\partial}{\partial t} |\Phi_t\rangle = H_{tot} |\Phi_t\rangle$ will then be fulfilled, and, accordingly, the evolution of the system with time will be correctly described. It is also easy to see that one-time operators $\tilde{C}^+$ and $\tilde{C}$ satisfy the conventional commutation relationships for fermions.

In the first order of the perturbation theory with respect to parameters $\mathbf{A}/T$ and $E_i^{\mathbf{p}, \mathbf{k}}/T$, it is possible to derive the following expressions:

$$\begin{aligned} \tilde{C}_{\sigma, \mathbf{p}}^+(t) &= \sum_{\mathbf{x}} \phi_{\mathbf{x},t}^{\mathbf{p}} C_{\sigma, \mathbf{x}}^+ + \sum_{\mathbf{x}, \mathbf{y}} \sum_{\mathbf{k}, \mathbf{z}} \left(\psi_{\mathbf{x}, \mathbf{y}, t}^{1, \mathbf{p}, \mathbf{k}} - \phi_{\mathbf{x}, t}^{\mathbf{p}} \phi_{\mathbf{y}, t}^{\mathbf{k}} \right) \phi_{\mathbf{z}, t}^{\mathbf{k}} C_{\sigma, \mathbf{x}}^+ C_{\sigma, \mathbf{y}}^+ C_{\sigma, \mathbf{z}}, \\ &+ \sum_{\mathbf{x}, \mathbf{y}} \sum_{\mathbf{k}, \mathbf{z}} \left(\psi_{\mathbf{x}, \mathbf{y}, t}^{0, \mathbf{p}, \mathbf{k}} + \psi_{\mathbf{x}, \mathbf{y}, t}^{1, \mathbf{p}, \mathbf{k}} - \phi_{\mathbf{x}, t}^{\mathbf{p}} \phi_{\mathbf{y}, t}^{\mathbf{k}} \right) \phi_{\mathbf{z}, t}^{\mathbf{k}} C_{\sigma, \mathbf{x}}^+ C_{-\sigma, \mathbf{y}}^+ C_{-\sigma, \mathbf{z}}, \\ \tilde{C}_{\sigma, \mathbf{p}}(t) &= \sum_{\mathbf{x}} \phi_{\mathbf{x}, t}^{* \mathbf{p}} C_{\sigma, \mathbf{x}} + \sum_{\mathbf{x}, \mathbf{y}} \sum_{\mathbf{k}, \mathbf{z}} \left(\psi_{\mathbf{x}, \mathbf{y}, t}^{* 1, \mathbf{p}, \mathbf{k}} - \phi_{\mathbf{x}, t}^{* \mathbf{p}} \phi_{\mathbf{y}, t}^{* \mathbf{k}} \right) \phi_{\mathbf{z}, t}^{\mathbf{k}} C_{\sigma, \mathbf{z}}^+ C_{\sigma, \mathbf{y}} C_{\sigma, \mathbf{x}}, \\ &+ \sum_{\mathbf{x}, \mathbf{y}} \sum_{\mathbf{k}, \mathbf{z}} \left(\psi_{\mathbf{x}, \mathbf{y}, t}^{* 0, \mathbf{p}, \mathbf{k}} + \psi_{\mathbf{x}, \mathbf{y}, t}^{* 1, \mathbf{p}, \mathbf{k}} - \phi_{\mathbf{x}, t}^{* \mathbf{p}} \phi_{\mathbf{y}, t}^{* \mathbf{k}} \right) \phi_{\mathbf{z}, t}^{\mathbf{k}} C_{-\sigma, \mathbf{z}}^+ C_{-\sigma, \mathbf{y}} C_{\sigma, \mathbf{x}}, \end{aligned} \quad (24)$$

whereas for ϕ and ψ we assume the fulfillment of the following relationships:

$$\begin{aligned} i \sum_{\mathbf{x}} \frac{\partial \phi_{\mathbf{x}, t}^{\mathbf{p}}}{\partial t} C_{\sigma, \mathbf{x}}^+ |0\rangle &= H_{tot} \sum_{\mathbf{x}} \phi_{\mathbf{x}, t}^{\mathbf{p}} C_{\sigma, \mathbf{x}}^+ |0\rangle, \\ i \sum_{\mathbf{x}, \mathbf{y}} \frac{\partial \psi_{\mathbf{x}, \mathbf{y}, t}^{0, \mathbf{p}, \mathbf{k}}}{\partial t} (C_{\sigma, \mathbf{x}}^+ C_{-\sigma, \mathbf{y}}^+ - C_{-\sigma, \mathbf{x}}^+ C_{\sigma, \mathbf{y}}^+) |0\rangle &= H_{tot} \sum_{\mathbf{x}, \mathbf{y}} \psi_{\mathbf{x}, \mathbf{y}, t}^{0, \mathbf{p}, \mathbf{k}} (C_{\sigma, \mathbf{x}}^+ C_{-\sigma, \mathbf{y}}^+ - C_{-\sigma, \mathbf{x}}^+ C_{\sigma, \mathbf{y}}^+) |0\rangle, \\ i \sum_{\mathbf{x}, \mathbf{y}} \frac{\partial \psi_{\mathbf{x}, \mathbf{y}, t}^{1, \mathbf{p}, \mathbf{k}}}{\partial t} C_{\sigma, \mathbf{x}}^+ C_{\sigma, \mathbf{y}}^+ |0\rangle &= H_{tot} \sum_{\mathbf{x}, \mathbf{y}} \psi_{\mathbf{x}, \mathbf{y}, t}^{1, \mathbf{p}, \mathbf{k}} C_{\sigma, \mathbf{x}}^+ C_{\sigma, \mathbf{y}}^+ |0\rangle, \end{aligned} \quad (25)$$

Assuming that it is possible to apply the perturbation theory to solving Eqs. (25) with accuracy up to linear terms with respect to $\mathbf{A}$, we may expect the fulfillment of relationships:

$$\phi_{\mathbf{x}, t}^{\mathbf{p}} = \delta_{\mathbf{x}}^{\mathbf{p}} e^{-iT_{\mathbf{p}} t},$$

$$\begin{aligned}\psi_{\mathbf{x},\mathbf{y},t}^{0,\mathbf{p},\mathbf{k}} &= \frac{1}{2} (\delta_{\mathbf{x}}^{\mathbf{p}} \delta_{\mathbf{y}}^{\mathbf{k}} + \delta_{\mathbf{y}}^{\mathbf{p}} \delta_{\mathbf{x}}^{\mathbf{k}}) e^{-i(T_{\mathbf{p}}+T_{\mathbf{k}}+E_0^{\mathbf{p},\mathbf{k}})t}, \\ \psi_{\mathbf{x},\mathbf{y},t}^{1,\mathbf{p},\mathbf{k}} &= \frac{1}{2} (\delta_{\mathbf{x}}^{\mathbf{p}} \delta_{\mathbf{y}}^{\mathbf{k}} - \delta_{\mathbf{y}}^{\mathbf{p}} \delta_{\mathbf{x}}^{\mathbf{k}}) e^{-i(T_{\mathbf{p}}+T_{\mathbf{k}}+E_1^{\mathbf{p},\mathbf{k}})t}.\end{aligned}\quad (26)$$

Now, the idea proper of calculations consists in the following. If we express the density of current $\mathbf{j}$ in operators $\tilde{C}^+$ and $\tilde{C}$, then it will be simple to calculate the mean value $\langle \mathbf{j} \rangle$. For this purpose, we will define the transformation which is reciprocal to that, given in Eq. (24):

$$\begin{aligned}C_{\sigma,\mathbf{x}}^+ &= \sum_{\mathbf{p}} \phi_{\mathbf{x},t}^{*\mathbf{p}} \tilde{C}_{\sigma,\mathbf{p}}^+ \\ &+ \sum_{\mathbf{p},\mathbf{k}} \sum_{\mathbf{y},\mathbf{z}} \left(\psi_{\mathbf{x},\mathbf{y},t}^{*1,\mathbf{p},\mathbf{k}} - \phi_{\mathbf{x},t}^{*\mathbf{p}} \phi_{\mathbf{y},t}^{*\mathbf{k}} \right) \phi_{\mathbf{y},t}^{\mathbf{z}} \tilde{C}_{\sigma,\mathbf{p}}^+ \tilde{C}_{\sigma,\mathbf{k}}^+ \tilde{C}_{\sigma,\mathbf{z}} \\ &+ \sum_{\mathbf{p},\mathbf{k}} \sum_{\mathbf{y},\mathbf{z}} \left(\psi_{\mathbf{x},\mathbf{y},t}^{*0,\mathbf{p},\mathbf{k}} + \psi_{\mathbf{x},\mathbf{y},t}^{*1,\mathbf{p},\mathbf{k}} - \phi_{\mathbf{x},t}^{*\mathbf{p}} \phi_{\mathbf{y},t}^{*\mathbf{k}} \right) \phi_{\mathbf{y},t}^{\mathbf{z}} \tilde{C}_{\sigma,\mathbf{p}}^+ \tilde{C}_{-\sigma,\mathbf{k}}^+ \tilde{C}_{-\sigma,\mathbf{z}}, \\ C_{\sigma,\mathbf{x}} &= \sum_{\mathbf{p}} \phi_{\mathbf{x},t}^{\mathbf{p}} \tilde{C}_{\sigma,\mathbf{p}} \\ &+ \sum_{\mathbf{p},\mathbf{k}} \sum_{\mathbf{y},\mathbf{z}} \left(\psi_{\mathbf{x},\mathbf{y},t}^{1,\mathbf{p},\mathbf{k}} - \phi_{\mathbf{x},t}^{\mathbf{p}} \phi_{\mathbf{y},t}^{\mathbf{k}} \right) \phi_{\mathbf{y},t}^{*\mathbf{z}} \tilde{C}_{\sigma,\mathbf{z}}^+ \tilde{C}_{\sigma,\mathbf{k}} \tilde{C}_{\sigma,\mathbf{p}} \\ &+ \sum_{\mathbf{p},\mathbf{k}} \sum_{\mathbf{y},\mathbf{z}} \left(\psi_{\mathbf{x},\mathbf{y},t}^{0,\mathbf{p},\mathbf{k}} + \psi_{\mathbf{x},\mathbf{y},t}^{1,\mathbf{p},\mathbf{k}} - \phi_{\mathbf{x},t}^{\mathbf{p}} \phi_{\mathbf{y},t}^{\mathbf{k}} \right) \phi_{\mathbf{y},t}^{*\mathbf{z}} \tilde{C}_{-\sigma,\mathbf{z}}^+ \tilde{C}_{-\sigma,\mathbf{k}} \tilde{C}_{\sigma,\mathbf{p}},\end{aligned}\quad (27)$$

Then, in the operators $\tilde{C}^+$ and $\tilde{C}$ the expression $c_{\sigma,\mathbf{k}}^+ c_{\sigma,\mathbf{k}+1}$, included in the expression for current density will be as follows:

$$\begin{aligned}c_{\sigma,\mathbf{p}}^+ c_{\sigma,\mathbf{p}+1} &= \sum_{\mathbf{p}',\mathbf{p}''} \phi_{\mathbf{p},t}^{*\mathbf{p}'} \phi_{\mathbf{p}+1,t}^{\mathbf{p}''} \tilde{C}_{\sigma,\mathbf{p}'}^+ \tilde{C}_{\sigma,\mathbf{p}''} \\ &- \sum_{\nu,\mathbf{k}',\mathbf{p}',\mathbf{p}''} \phi_{\mathbf{p},t}^{*\mathbf{p}'} \phi_{\mathbf{p}+1,t}^{\mathbf{p}''} \tilde{C}_{\sigma,\mathbf{p}'}^+ \tilde{C}_{\nu,\mathbf{k}'}^+ \tilde{C}_{\nu,\mathbf{k}''} \tilde{C}_{\sigma,\mathbf{p}''} \\ &+ \sum_{\mathbf{k},\mathbf{p},\mathbf{z}} \sum_{\mathbf{y}} \chi_{1,\mathbf{z}+\mathbf{y}}^{*\mathbf{k},\mathbf{p}} \chi_{1,\mathbf{y}}^{\mathbf{k}+\mathbf{z},\mathbf{p}-\mathbf{z}+1} \sum_{\nu,\mathbf{k}',\mathbf{k}''} \sum_{\mathbf{p}',\mathbf{p}''} \psi_{\mathbf{p},\mathbf{k},t}^{*1,\mathbf{p}',\mathbf{k}'} \psi_{\mathbf{p}-\mathbf{z}+1,\mathbf{k}+\mathbf{z},t}^{1,\mathbf{p}'',\mathbf{k}''} \tilde{C}_{\sigma,\mathbf{p}'}^+ \tilde{C}_{\nu,\mathbf{k}'}^+ \tilde{C}_{\nu,\mathbf{k}''} \tilde{C}_{\sigma,\mathbf{p}''} \\ &+ \sum_{\mathbf{k},\mathbf{p},\mathbf{z}} \sum_{\mathbf{y}} \chi_{1,\mathbf{z}+\mathbf{y}}^{*\mathbf{k},\mathbf{p}} \chi_{1,\mathbf{y}}^{\mathbf{k}+\mathbf{z},\mathbf{p}-\mathbf{z}+1} \sum_{\nu,\mathbf{k}',\mathbf{k}''} \sum_{\mathbf{p}',\mathbf{p}''} \psi_{\mathbf{p},\mathbf{k},t}^{*0,\mathbf{p}',\mathbf{k}'} \psi_{\mathbf{p}-\mathbf{z}+1,\mathbf{k}+\mathbf{z},t}^{0,\mathbf{p}'',\mathbf{k}''} \tilde{C}_{\sigma,\mathbf{p}'}^+ \tilde{C}_{-\sigma,\mathbf{k}'}^+ \tilde{C}_{-\sigma,\mathbf{k}''} \tilde{C}_{\sigma,\mathbf{p}''} \\ &+ \sum_{\mathbf{k},\mathbf{p},\mathbf{z}} \sum_{\mathbf{y}} \chi_{0,\mathbf{z}+\mathbf{y}}^{*\mathbf{k},\mathbf{p}} \chi_{0,\mathbf{y}}^{\mathbf{k}+\mathbf{z},\mathbf{p}-\mathbf{z}+1} \sum_{\nu,\mathbf{k}',\mathbf{k}''} \sum_{\mathbf{p}',\mathbf{p}''} \psi_{\mathbf{p},\mathbf{k},t}^{*1,\mathbf{p}',\mathbf{k}'} \psi_{\mathbf{p}-\mathbf{z}+1,\mathbf{k}+\mathbf{z},t}^{1,\mathbf{p}'',\mathbf{k}''} \tilde{C}_{\sigma,\mathbf{p}'}^+ \tilde{C}_{-\sigma,\mathbf{k}'}^+ \tilde{C}_{-\sigma,\mathbf{k}''} \tilde{C}_{\mu,\mathbf{p}''} \\ &+ \sum_{\mathbf{k},\mathbf{p},\mathbf{z}} \sum_{\mathbf{y}} \chi_{0,\mathbf{z}+\mathbf{y}}^{*\mathbf{k},\mathbf{p}} \chi_{0,\mathbf{y}}^{\mathbf{k}+\mathbf{z},\mathbf{p}-\mathbf{z}+1} \sum_{\nu,\mathbf{k}',\mathbf{k}''} \sum_{\mathbf{p}',\mathbf{p}''} \psi_{\mathbf{p},\mathbf{k},t}^{*0,\mathbf{p}',\mathbf{k}'} \psi_{\mathbf{p}-\mathbf{z}+1,\mathbf{k}+\mathbf{z},t}^{0,\mathbf{p}'',\mathbf{k}''} \tilde{C}_{\sigma,\mathbf{p}'}^+ \tilde{C}_{-\sigma,\mathbf{k}'}^+ \tilde{C}_{-\sigma,\mathbf{k}''} \tilde{C}_{\sigma,\mathbf{p}''}\end{aligned}\quad (28)$$

Limiting ourselves only to unrelativistic terms in the equation for $\frac{\partial \mathbf{j}_{\mathbf{q}}}{\partial t}$ with an accuracy of up to the second order with respect to parameter $\mathbf{A}$, it is possible to derive

$$\left\langle \frac{\partial \mathbf{j}_{\mathbf{q}}}{\partial t} \right\rangle = -\frac{e^2 L^{-3}}{mc} \sum_{\sigma,\mathbf{p},\mathbf{l}} \frac{\partial \mathbf{A}_{\mathbf{q}-\mathbf{l}}}{\partial t} \langle c_{\sigma,\mathbf{p}}^+ c_{\sigma,\mathbf{p}+1} \rangle = -\frac{e^2 n}{mc} \frac{\partial \mathbf{A}_{\mathbf{q}}}{\partial t}, \quad (29)$$

or

$$\langle \mathbf{j}_{\mathbf{q}} \rangle = -\frac{e^2 n}{mc} \mathbf{A}_{\mathbf{q}}, \quad (30)$$

which just constitutes the essential point of Londons' hypothesis.

IV. CONCLUSIONS

In our opinion, it is possible to subdivide the existing, relevant from the viewpoint of physics studies on the superconductivity theory into two categories enumerated below.

1. Systems with Bose–Einstein Condensation

In such works, the Hamiltonians with electron-electron attraction are introduced, which are considered as initial ones. In this, the boson operators corresponding to the bound states of electron pairs are used to determine the ground state of the system.

The studies devoted to dealing with this problem are, as a whole, not systematized. Therefore, it is very difficult to single out the most significant of them. It should also be noted that some of the papers anticipate the BCS theory⁹. Therefore, the use of the term "Cooper pairs" seems to be not quite suitable, especially from the viewpoint of the remarks, given in the introduction to our paper.

With some provision, the modifications of the popular $t - j$ model may be related to those studies. With a certain selection of parameters of the model, for example $j > t$, a possibility arises of the formation of bound pairs. The condition $j > t$ is linked with a short-wave character of the interaction in the Hubbard form and Heisenberg's uncertainty principle. The authors of these works themselves call the choice of such parameters as "unphysical"¹⁰.

Now, the introduction of Hamiltonian to the studies devoted to the Bose condensation in Fermi gas is either unsubstantiated or causes skepticism in the minds of most physicists¹¹. In this sense, we may reckon the authors of these works are so far investigating hypothetical objects.

The concept of Bose condensate in such systems has a distinctly expressed sense. That is why the Ginzburg-Landau theory using the wave function of Bose condensate in the definition of the order parameter of the system is quite valid.

2. Systems with the Fermi Condensation

In this case, the electron-electron attraction arises only with an effective potential which is calculated in an assumption that the wave function of the ground state coincides in its form with the function of the ground state of a normal metal $|\Phi_0\rangle = \prod_{\sigma, k < K_F} C_{\sigma, \mathbf{k}}^+ |0\rangle$. The question to be settled resides merely in selecting the operators $C_{\sigma, \mathbf{k}}^+$. The effective potential is calculated in the framework of refining operators $C_{\sigma, \mathbf{k}}^+$ when solving the problem of renormalization and accounting for electron-electron correlations. From this viewpoint, the operators $C_{\sigma, \mathbf{k}}^+$ are linked in a self-consistent manner with the effective potential of the system, and there is no additional problem involving the definition of the wave function of the ground state.

The substantiation of a possibility of the origination of the effective electron-electron attraction in metals is given in study⁷.

There is, presumably, so far no phenomenological theory available on the systems of such a kind.

Acknowledgments

We are grateful to Prof. A.A. Rukhadze for being helpful in the statement of the problem.

APPENDIX

Taking into account Eqs. (12), it is possible to expect the fulfillment of the following conditions:

$$\begin{aligned}
 \chi_{0, \mathbf{q}}^{\mathbf{p}, \mathbf{k}} &= \chi_{0, \mathbf{k}-\mathbf{p}-\mathbf{q}}^{\mathbf{p}, \mathbf{k}}, \chi_{0, \mathbf{q}}^{\mathbf{p}, \mathbf{k}} = \chi_{0, -\mathbf{q}}^{\mathbf{k}, \mathbf{p}}, \\
 \chi_{1, \mathbf{q}}^{\mathbf{p}, \mathbf{k}} &= -\chi_{1, \mathbf{k}-\mathbf{p}-\mathbf{q}}^{\mathbf{p}, \mathbf{k}}, \chi_{1, \mathbf{q}}^{\mathbf{p}, \mathbf{k}} = \chi_{1, -\mathbf{q}}^{\mathbf{k}, \mathbf{p}}; \\
 \sum_{\mathbf{z}} \chi_{0, \mathbf{z}}^{\mathbf{p}, \mathbf{k}+\mathbf{q}} \chi_{0, \mathbf{q}-\mathbf{z}}^{\mathbf{k}, \mathbf{p}+\mathbf{q}} &= \frac{1}{2} (\delta_{\mathbf{q}}^0 + \delta_{\mathbf{p}}^{\mathbf{k}}) \\
 \sum_{\mathbf{z}} \chi_{1, \mathbf{z}}^{\mathbf{p}, \mathbf{k}+\mathbf{q}} \chi_{1, \mathbf{q}-\mathbf{z}}^{\mathbf{k}, \mathbf{p}+\mathbf{q}} &= \frac{1}{2} (\delta_{\mathbf{q}}^0 - \delta_{\mathbf{p}}^{\mathbf{k}}).
 \end{aligned} \tag{A.1}$$

The fulfillment of these conditions is conducive to that the fermions commutation relationships for operators C^+ and C , as well, as their reciprocal transformation (13), are violated only beginning with 4-linear terms on the operators of creation and annihilation. It is just this circumstance that is employed in constructing the model Hamiltonian of the system.

* Electronic address: yurinoffice@aport2000.ru

¹ J. Bardeen, L. N. Cooper, and J. R. Schrieffer, *Phys. Rev.* **106**, 162 (1957); **108**, 1175 (1957).

² Quantum Mechanics, Harry J. Lipkin, American Elsevier publishing Co., Inc., New York, 1973.

- ³ A. O. E. Animalu, Intermediate Quantum Theory of Crystalline Solids, Prentice-Hall, Inc., Englewood Cliffs, New Jersey, 1977.
- ⁴ G. M. Eliashberg, *JETP* **11**, 696, (1960); **12**, 1000 (1961).
- ⁵ E. G. Maksimov, D. Yu. Savrasov, S. Yu. Savrasov, *Usp. Fiz. Nauk (Rus)***40** (4), 337 (1997).
- ⁶ High-Temperature Superconductivity, Eds. V.L.Ginzburg, D.A.Kirzhnits, New York: Consultant Bureau, 1982.
- ⁷ I. M. Yurin, V. B. Kalinin, *cond-mat/0102407*; *Appl.Phys.(Rus)* **2**, 12 (2003).
- ⁸ I. M. Yurin, V. B. Kalinin, *cond-mat/0202486*.
- ⁹ M. R. Schafroth, *Phys. Rev.* **96**, 1442 (1954).
- ¹⁰ L. P. Pryadko, *cond-mat/0306342*.
- ¹¹ M. R. Norman, *J. Phys. Chem. Sol.* **54**, 1165 (1993).